



Certificate of Completion Tuberculosis Risk Assessment and/or Examination

To satisfy **job-related requirements** in the California Education Code, Sections 49406 and 87408.6 and the California Health and Safety Code, Sections 1597.055, 121525, 121545 and 121555.

First and Last Name of the person assessed and/or examined:
Date of assessment and/or examination:mo./day/yr.
Date of Birth:mo./day/yr.
The above named patient has submitted to a tuberculosis risk assessment. The patient does not have risk factors, or if tuberculosis risk factors were identified, the patient has been examined and determined to be free of infectious tuberculosis.
X
Please print, place label or stamp with Health Care Provider Name and Address (include Number, Street, City, State, and Zip Code):
Telephone and FAX:



School Staff and Volunteers: Tuberculosis Risk Assessment



Job-related requirement for child care, pre-K, K-12, and community colleges

The purpose of this tool is to identify **adults** with infectious tuberculosis (TB) to prevent them from spreading TB. Use of this risk assessment is required in the California Education Code, Sections 49406 and 87408.6 and the California Health and Safety Code, Sections 1597.055 and 121525, 121545, and 121555.

The law requires that a health care provider administer this risk assessment. A health care provider, as defined for this purpose, is any organization, facility, institution or person licensed, certified or otherwise authorized or permitted by state law to deliver or furnish health services. Any person administering this risk assessment is to have training in the purpose and significance of the risk assessment and Certificate of Completion.

Name of Employee/Volunteer Assessed for TB Risk Factors:		
Assessment Date:	Date of Birth:	
History of Tuberculosis I	nfection or Disease (Check appropriate box below)	
chest x-ray (if none performed in pre- physician assistant, or nurse practition has been followed by an x-ray, and w	ositive TB test (infection) or TB disease, then a symptom review and vious 6 months) should be performed at initial hire by a physician, oner. Once a person has a documented positive test for TB infection that was determined to be free of infectious TB, the TB risk assessment (and If an employee or volunteer becomes symptomatic for TB, then he/she care provider.	
☐ No (Assess for Risk Factors for Tube	erculosis using box below)	
Release Assay (IGRA). Re-testing with T negative, and have new risk factors since chest x-ray, and if normal, treatment for T Latent Tuberculosis Infection: A Guide for Prin	perform a Mantoux tuberculin skin test (TST) or Interferon Gamma ST or IGRA should only be done in persons who previously tested the last assessment. A positive TST or IGRA should be followed by a B infection considered. (Centers for Disease Control and Prevention [CDC]. mary Health Care Providers. 2013) of TB: prolonged cough, coughing up blood, fever, night sweats, weight	
	a TST or IGRA, chest x-ray, symptom screen, and if indicated, sputumes and nucleic acid amplification testing. A negative TST or IGRA does	
Close contact to someone with infection.	ctious TB disease at any time	
Foreign-born person from a country Includes any country other than the Uni Europe. IGRA is preferred over TST for	ited States, Canada, Australia, New Zealand, or a country in western or northern	
AN A	ed States, Canada, Australia, New Zealand, or a country in western or northern	
☐ Volunteered, worked or lived in a cor	rectional or homeless facility	